arXiv:math/0201129v1 [math.AG] 15 Jan 2002

DIMENSIONS OF JET SCHEMES OF LOG SINGULARITIES

TAKEHIKO YASUDA

Abstract. We characterize Kawamata log terminal singularities and log canonical singularities by dimensions of jet schemes. Our main result is Theorem 2.4.

Introduction

For birational geometry, it is natural to consider a pair (X, D) of a normal variety X and a $\mathbb{Q}$-divisor on it, and to consider singularities of pairs. *KLT (Kawamata log terminal)* and *LC (log canonical)* singularities form important classes of log singularities. They are defined by using a log-resolution and discrepancies. When X is $\mathbb{Q}$-Gorenstein, we can naturally define KLT and LC also for a pair (X, qY) where Y is a closed subscheme of X and $q \in \mathbb{Q}_{>0}$.

In [Mus01], Mustaţă proposed a new point of view in the study of log singularities. He characterized KLT and LC pairs (X, qY) with X smooth via dimensions of jet schemes of Y. The aim of this short paper is to extend his result to the case X is $\mathbb{Q}$-Gorenstein and to shorten his proof. As he did, we also use the motivic integration as a main tool, which is invented by Kontsevich [Kon95] and extended to singular varieties by Denef and Loeser [DL99a].

Acknowledgements. This paper forms a part of my master's thesis. First, I would like to thank Yujiro Kawamata for his encouragement and advise. I also thank Nobuyuki Kakimi for useful conversation.

1. Motivic integration

In this section, we review the motivic integration invented by Kontsevich [Kon95] and extended by Denef, Loeser [DL99a]. Craw's paper [Cra99] is a nice introduction.

Let k be an algebraically closed field of characteristic zero.

1.1. Extended Grothendieck rings of varieties.

The *Grothendieck ring of k-varieties*, denoted $K_0(\mathrm{Var}/k)$, is the abelian group generated by the isomorphism classes $[X]$ of k-varieties with the relations $[X] = [X \setminus Y] + [Y]$ if Y is a closed subvariety of X. The ring structure is defined by $[X][Y] =$

$[X \times Y]$. Let $\mathbb{L}$ be the class $[\mathbb{A}^1]$ of the affine line and let $\mathbb{M}$ be the localization $K_0(\mathrm{Var}/k)[\mathbb{L}^{-1}]$. We define the *dimension* of an element of $\mathbb{M}$ by

$$\dim(\sum_i [X_i]\mathbb{L}^{m_i}) := \max_i(\dim X_i + m_i),$$

with the convention $\dim \emptyset := -\infty$. Then, the following hold: for $\alpha,\ \beta \in \mathbb{M}$,

- $\dim \alpha\beta = \dim \alpha + \dim \beta$,
- $\dim(\alpha + \beta) \le \max\{\dim \alpha, \dim \beta\}$ and the equality holds if $\dim \alpha \ne \dim \beta$.

If we set $F_m\mathbb{M} := \{\alpha \in \mathbb{M} | \dim \alpha \le m\}$, then $(F_m\mathbb{M})_m$ is an ascending filtration with $F_m\mathbb{M} \cdot F_n\mathbb{M} \subset F_{m+n}\mathbb{M}$. We define the *complete Grothendieck ring of k-varieties* by

$$\hat{\mathbb{M}} := \varprojlim \mathbb{M}/F_m\mathbb{M} \quad (m \to -\infty).$$

Any element α of $\hat{\mathbb{M}}$ is expressed as $\alpha = \sum_{m\in\mathbb{Z}} \alpha_m$ such that

- $\alpha_m = 0$ or $\dim \alpha_m = m$,
- $\alpha_m = 0$ for $m \gg 0$.

Then we define the *dimension* of α to be $\max\{m|\alpha_m \ne 0\}$.

We denote by $\hat{\mathbb{M}}^{\mathbb{Q}}$ the ring $\hat{\mathbb{M}}[\mathbb{L}^q; q \in \mathbb{Q}]$. Any element of $\hat{\mathbb{M}}^{\mathbb{Q}}$ is expressed as $\alpha\mathbb{L}^q$ with $\alpha \in \hat{\mathbb{M}},\ q \in \mathbb{Q}$. We define the *dimension* of $\alpha\mathbb{L}^q$ to be $\dim \alpha + q$.

1.2. **Jet schemes and motivic measures.**

Let X be a k-scheme. For $n \in \mathbb{Z}_{\ge 0} \cup \{\infty\}$, an *$n$-jet* on X is a morphism

$$\mathrm{Spec}\, k[[t]]/(t^{n+1}) \to X,$$

where we used the convention $(t^\infty) = (0) \subset k[[t]]$. The moduli scheme of the n-jets on X always exists. We call it the *n-jet scheme* of X and denote it by $L_n(X)$. If X is of finite type, then for $n < \infty$, so is $L_n(X)$. If X is smooth and of pure dimension d, then, for each $n \in \mathbb{Z}_{\ge 0}$, the natural projection $L_{n+1}(X) \to L_n(X)$ is a Zariski locally trivial $\mathbb{A}^d$-bundle.

Now assume that X is a k-variety of pure dimension d. Let $\pi_n : L_\infty(X) \to L_n(X)$ be the canonical projection.

Definition 1.1. A subset A of $L_\infty(X)$ is *stable at level n* if we have:

1. $\pi_n(A)$ is a constructible subset of $L_n(X)$,
2. $A = \pi_n^{-1}\pi_n(A)$,
3. for any $m \ge n$, the projection $\pi_{m+1}(A) \to \pi_m(A)$ is a piecewise trivial $\mathbb{A}^d$-bundle.

(A morphism $f : Y \to X$ of schemes is called a *piecewise trivial $\mathbb{A}^d$-bundle* if there is a stratification $X = \coprod X_i$ such that $f|_{f^{-1}(X_i)} : f^{-1}(X_i) \to X_i$ is isomorphic to $X_i \times \mathbb{A}^d \to X_i$ for each i.) A subset A of $L_\infty(X)$ is *stable* if it is stable at level n for some $n \in \mathbb{Z}_{\ge 0}$.

We see that the stable subsets of $L_\infty(X)$ constitute a Boolean algebra, that is, stable under finite intersection and finite union. For a stable subset

A of $L_\infty(X)$, an element $[\pi_n(A)]\mathbb{L}^{-nd} \in \hat{\mathbb{M}}$ is independent of the choice of $n \gg 0$. (We can define the class of a constructible set in $\mathbb{M}$ and $\hat{\mathbb{M}}$ in the evident fashion.) So the map

$$\begin{aligned} \mu_X : \{\text{stable subsets of } L_\infty(X)\} &\to \hat{\mathbb{M}} \\ A \qquad\qquad &\mapsto [\pi_n(A)]\mathbb{L}^{-nd}\ (n \gg 0) \end{aligned}$$

is a finite additive measure. We can extend μ_X to the family of the *measurable subsets* of $L_\infty X$, which is a family 'big enough'. For details, see [DL99b], [Loo00]. We call μ_X the *motivic measure* on $L_\infty X$.

Let $A \subset L_\infty(X)$ be a measurable subset and $\nu : A \to \mathbb{Q} \cup \{\infty\}$ a function. We say that ν is a *measurable function* if the fibers are measurable and $\mu_X(\nu^{-1}(\infty)) = 0$.

Definition 1.2. For a measurable function ν, we formally define the *motivic integration* of $\mathbb{L}^\nu$ by

$$\int_A \mathbb{L}^\nu d\mu_X := \sum_{n \in \mathbb{Q}} \mu_X(\nu^{-1}(n))\mathbb{L}^n.$$

We say that $\mathbb{L}^\nu$ is *integrable* if this infinite sum converges in $\hat{\mathbb{M}}^{\mathbb{Q}}$.

Let $Y \subset X$ be a closed subscheme and $\mathfrak{a}$ its ideal sheaf. A closed point $\gamma \in L_\infty(X)$ corresponds to a morphism $\gamma' : \operatorname{Spec} k[[t]] \to X$. The function

$$\begin{aligned} F_Y : L_\infty(X) &\to \mathbb{Z}_{\geq 0} \cup \{\infty\} \\ \gamma\ &\mapsto n \ \text{ if } (\gamma')^{-1}\mathfrak{a} = (t^n) \end{aligned}$$

is a measurable function. For a $\mathbb{Q}$-divisor $D = \sum_i q_i D_i$ on X with D_i a prime divisor, we define a measurable function $F_D := \sum_i q_i F_{D_i}$. The following *explicit formula* give a way to compute motivic integrations.

Lemma 1.3. *Assume that X is smooth and $\sum_{i=1}^s D_i$ is a SNC divisor on X with D_i a prime divisor.*

Let $m_i \in \mathbb{Z}_{\geq 0}$, $(1 \leq i \leq s)$, and put $J := \{i | m_i > 0\} \subset \{1, \ldots, s\}$. Then we have

$$\mu_X(\bigcap_i F_{D_i}^{-1}(m_i)) = [D_J^\circ](\mathbb{L} - 1)^{|J|}\mathbb{L}^{-\sum m_i},$$

where

$$D_J^\circ := \bigcap_{i \in J} D_i \Big\backslash \bigcup_{i \in \{1,\ldots,s\} \setminus J} D_i.$$

Proof. See [Cra99, the proof of Thm. 2.15]. □

2. MAIN THEOREM

Let X be a normal variety of dimension d. Assume that X is $\mathbb{Q}$-Gorenstein, that is, for some $r \in \mathbb{Z}_{>0}$, rK_X is a Cartier divisor. For a resolution $p : \tilde{X} \to X$, the *relative canonical divisor* $K_{\tilde{X}/X}$ is

$$K_{\tilde{X}/X} := \frac{1}{r}(rK_{\tilde{X}} - p^*(rK_X)).$$

Let Y be a closed subscheme of X and $\mathfrak{a}$ its ideal sheaf. By Hironaka's theorem, there are a resolution $p : \tilde{X} \to X$ and a SNC divisor $\sum_i D_i$ on $\tilde{X}$ such that

- $p^{-1}\mathfrak{a} = \mathcal{O}_{\tilde{X}}(-\sum_i y_i D_i)$ for some $y_i \in \mathbb{Z}_{\geq 0}$,
- $K := K_{\tilde{X}/X} = \sum_i a_i D_i$ for some $a_i \in \mathbb{Q}$.

Fix these notations through the rest of the paper. In the proof of Theorem 2.4, we suppose additional conditions on p.

Definition 2.1. For $q \in \mathbb{Q}_{>0}$, we say that the pair (X, qY) is *KLT (Kawamata log terminal)*, resp. *LC (log canonical)* if for every i, $-qy_i + a_i + 1 > 0$, resp. $-qy_i + a_i + 1 \geq 0$.

Let X_{reg} be the smooth locus of X and $\iota : X_{\mathrm{reg}} \hookrightarrow X$ the inclusion. Then $\omega_X^{[r]} = \iota_*((\Omega_X^d)^{\otimes r})$ is an invertible sheaf. We define an ideal sheaf $\mathcal{J} \subset \mathcal{O}_X$ by the following equation:

$$\mathcal{J}\omega_X^{[r]} = \mathrm{Image}((\Omega_X^d)^{\otimes r} \to \omega_X^{[r]}).$$

Let $Z \subset X$ be the closed subscheme associated to $\mathcal{J}$. Then $\mathrm{Supp}\, Z = X_{\mathrm{sing}}$. The following is a variation of the transformation rule, a key of the proof of the main theorem.

Theorem 2.2. *Let $A \subset L_\infty X$ be a measurable subset and $\nu : A \to \mathbb{Q} \cup \{\infty\}$ a measurable function. Then we have the following equality:*

$$\int_A \mathbb{L}^{\nu + (1/r)\cdot F_Z} d\mu_X = \int_{p_\infty^{-1}(A)} \mathbb{L}^{\nu \circ p_\infty - F_K} d\mu_{\tilde{X}}.$$

Proof. It is a direct consequence of the transformation rule [DL99a, Lem. 3.3]. □

For each $e \in \mathbb{Z}_{\geq 0}$, we put $A_e := F_Z^{-1}(e)$. For each $n, e \in \mathbb{Z}_{\geq 0}$, we define

$$L_n^e(Y) := L_n(Y) \cap \pi_n(A_e),$$

where we take the intersection in $L_n(X)$. Of course, $L_n^e(Y)$ depends on the inclusion $Y \subset X$.

Lemma 2.3. *There is a positive integer θ such that for any $n, e \in \mathbb{Z}_{\geq 0}$ with $n \geq \theta e$, the natural projection $\pi_{n+1}(A_e) \to \pi_n(A_e)$ is a piecewise trivial $\mathbb{A}^d$-bundle.*

Proof. See [DL99a, Lem. 4.1]. □

Theorem 2.4. *Suppose* $\operatorname{Supp} Z \supset X_{\mathrm{sing}}$. *Let* θ *be a positive integer as in Lemma 2.3. For* $l \in \mathbb{Z}_{>0}$, *let* lY *be the closed subscheme of* X *associated to the ideal sheaf* $\mathfrak{a}^l$. *Suppose that* $\mathfrak{a}^l \subset \mathcal{J}^\theta$. *Then,*

(1) (X, qY) *is KLT iff for any* $e, n \in \mathbb{Z}_{\geq 0}$ *with* $n \geq \theta e$,

$$\dim L_n^e(lY) + e/r < (n+1)(d - q/l).$$

(2) (X, qY) *is LC iff for any* $e, n \in \mathbb{Z}_{\geq 0}$ *with* $n \geq \theta e$,

$$\dim L_n^e(lY) + e/r \leq (n+1)(d - q/l).$$

Proof. We prove only (1). For each $n \in \mathbb{Z}_{\geq 0}$, let $B_n := F_Y^{-1}(n)$ and $B_{\geq n} := F_Y^{-1}(\mathbb{Z}_{\geq n})$. For each $e, n \in \mathbb{Z}_{\geq 0}$ with $n \geq \theta e$, consider the following element of $\hat{\mathbb{M}}^{\mathbb{Q}}$:

$$\begin{aligned} S(e,n) &:= \int_{B_{n+1} \cap A_e} \mathbb{L}^{qF_Y + (1/r)\cdot F_Z} d\mu_X \\ &= \mu_X(B_{n+1} \cap A_e) \mathbb{L}^{(n+1)q + e/r}. \end{aligned}$$

By Lemma 2.3,

$$\begin{aligned} S(e,n) =& (\mu_X(B_{\geq n+1} \cap A_e) - \mu_X(B_{\geq n+2}) \cap A_e)) \mathbb{L}^{(n+1)q+e/r} \\ =& ([L_n^e(Y)] - [L_{n+1}^e(Y)]\mathbb{L}^{-d}) \mathbb{L}^{-nd + (n+1)q + e/r}. \end{aligned}$$

By Lemma 2.3 again, we have $\dim L_{n+1}^e(Y) \leq \dim L_n^e(Y) + d$, and hence

$$\dim S(e,n) \leq \dim L_n^e(Y) - nd + (n+1)q + e/r. \tag{2.1}$$

If the equality does not hold, then $\dim L_{n+1}^e(Y) = \dim L_n^e(Y) + d$.

We suppose that $p^{-1}\mathcal{J} = \mathcal{O}_{\tilde{X}}(-\sum_i z_i D_i)$ with $z_i > 0$.

We set $\tilde{Y} := \sum_i y_i D_i$ and $\tilde{Z} := \sum_i z_i D_i$. From Theorem 2.2, we have

$$S(e,n) = \int \mathbb{L}^{qF_{\tilde{Y}} - F_K} d\mu_{\tilde{X}}$$

where the domain of the integration is

$$F_{\tilde{Y}}^{-1}(n+1) \cap F_{\tilde{Z}}^{-1}(e).$$

From Lemma 1.3, we obtain

$$S(e,n) = \sum_{J \subset \{1,\dots,s\}} \sum_{\mathbf{m} \in M} [D_J^\circ](\mathbb{L}-1)^{|J|} \mathbb{L}^{-\sum(-qy_i + a_i + 1)m_i},$$

where

$$\begin{aligned} M &= M(J,n,e) \\ &:= \{\mathbf{m} = (m_i)_{i \in J} \in (\mathbb{Z}_{>0})^J \mid \textstyle\sum y_i m_i = n \text{ and } \sum z_i m_i = e\}. \end{aligned}$$

For each J and $\mathbf{m}$, the highest term of

$$[D_J^\circ](\mathbb{L}-1)^{|J|} \mathbb{L}^{-\sum(-qy_i + a_i + 1)m_i}$$

has a positive coefficient. Hence,

$$\dim S(e,n) = \max_{\substack{J\subset\{1,\dots,s\}\\ \mathbf{m}\in M}} (d - \sum_i (-qy_i + a_i + 1)m_i). \tag{2.2}$$

'Only if' part: The proof is by contradiction. So assume that (X, qY) is KLT, that is, for every i, $-qy_i + a_i + 1 > 0$, and that for some $n', e' \in \mathbb{Z}_{\geq 0}$ with $n' > \theta e'$,

$$\dim L^{e'}_{n'}(Y) + e'/r \geq (n'+1)(d-q).$$

By (2.2), we have $\dim S(e', n') < d$. On the other hand, since

$$\dim L^{e'}_{n'}(Y) - n'd + (n'+1)q + e'/r \geq d,$$

the equality in (2.1) does not hold. Hence $\dim L^{e'}_{n'+1}(Y) = \dim L^{e'}_{n'}(Y) + d$ and hence $\dim L^{e'}_{n'+1}(Y) + e'/r \leq ((n'+1)+1)(d-q)$. By the same argument, we have $\dim L^{e'}_{n'+2}(Y) = \dim L^{e'}_{n'+1}(Y) + d$ and so on. It contradicts Lemma 2.5.

'If' part: From inequality (2.1), we have that for any e, n with $n \geq \theta e$,

$$\dim S(e,n) < d. \tag{2.3}$$

In view of (2.2) and (2.3), it is easy to find that for any i, $-qy_i + a_i + 1 > 0$, that is, (X, qY) is KLT. We have thus completed the proof. □

Lemma 2.5. *Then for each $e \in \mathbb{Z}_{\geq 0}$, there is a real numbers b such that $b < d$ and for every $n \in \mathbb{Z}_{\geq 0}$,*

$$\dim L^e_n(Y) \leq bn + (\text{const}).$$

Proof. It suffices to find an increasing linear function $\psi : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$ and a real number $b < d$ such that

$$\dim L^e_{\psi(n)}(Y) \leq b\psi(n) + (\text{const}).$$

First, the case Y reduced: For $m, n \in \mathbb{Z}_{\geq 0}$ with $m \geq n$, we denote by π^m_n the natural projection $L_m(X) \to L_n(X)$. By Greenberg's theorem [Gre66, Cor. 1], there is a linear function $g : \mathbb{Z}_{\geq 0} \to \mathbb{Z}_{\geq 0}$ such that

- for every $n \in \mathbb{Z}_{\geq 0}$, $g(n) \geq n$,
- $\pi_n(L_\infty(Y)) = \pi^{g(n)}_n(L_{g(n)}(Y))$.

Since $L^e_{g(n)}(Y) \subset (\pi^{g(n)}_n)^{-1}\pi^{g(n)}_n(L^e_{g(n)}(Y))$,

$$\dim L^e_{g(n)}(Y) \leq \dim(\pi^{g(n)}_n)^{-1}\pi^{g(n)}_n(L^e_{g(n)}(Y)).$$

By the definition of g,

$$\pi^{g(n)}_n(L^e_{g(n)}(Y)) = \pi_n(L^e_\infty(Y)).$$

Let d' be the dimension of Y. By [DL99a, Lem 4.3],

$$\dim \pi^{g(n)}_n(L^e_{g(n)}(Y)) = \dim \pi_n(L^e_\infty(Y)) \leq \dim(n+1)d'.$$

Hence, by Lemma 2.3,

$$\begin{aligned}\dim L^e_{g(n)}(Y) &\leq (n+1)d' + (g(n)-n)d + (\mathrm{const})\\ &= g(n)d + n(d'-d) + (\mathrm{const})\\ &= g(n)(d + (d'-d)/C_1) + (\mathrm{const}),\end{aligned}$$

where C_1 is the constant such that $g(n) = C_1 n + (\mathrm{const})$. We have thus proved the assertion in this case.

The general case: It suffices to show the case where $Y = l(Y_{\mathrm{red}})$ for some $l \in \mathbb{Z}_{>0}$. By the definitions, we have the following

$$\begin{aligned}L^e_{ln-1}(Y) &= \pi_{ln-1}(F_Y^{-1}(ln) \cap A_e),\\ L^e_{n-1}(Y_{\mathrm{red}}) &= \pi_{n-1}(F_{Y_{\mathrm{red}}}^{-1}(n) \cap A_e).\end{aligned}$$

Because for n, e with $n \geq \theta e$, $F_Y^{-1}(ln) \cap A_e = F_{Y_{\mathrm{red}}}^{-1}(n) \cap A_e$ is stable at level n, we have $L^e_{ln-1}(Y) = (\pi^{ln-1}_{n-1})^{-1} L^e_n(Y_{\mathrm{red}})$. Therefore, for some $b < d$,

$$\begin{aligned}\operatorname{codim}(L^e_{ln-1}(Y)/\pi_{ln-1}(A_e)) &= \operatorname{codim}(L^e_{n-1}(Y_{\mathrm{red}})/\pi_{n-1}(A_e))\\ &\geq (d-b)\varphi(n) + (\mathrm{const}).\end{aligned}$$

Because $\dim \pi_n(A_e) = dn + (\mathrm{const})$, we have

$$\begin{aligned}\dim L^e_{ln-1}(Y) &\leq dln - (d-b)n + (\mathrm{const})\\ &\leq (d - (d-b)/l)(ln-1) + (\mathrm{const}).\end{aligned}$$

This completes the proof. □

References

[Cra99] Alastair Craw. An introduction to motivic integration. preprint, `math.AG/9911179`, 1999.

[DL99a] Jan Denef and François Loeser. Germs of arcs on singular algebraic varieties and motivic integration. *Invent. Math.*, 135(1):201–232, 1999.

[DL99b] Jan Denef and François Loeser. Motivic integration, quotient singularities and the McKay correspondence. preprint, `math.AG/9903187`, 1999.

[Gre66] Marvin J. Greenberg. Rational points in Henselian discrete valuation rings. *Publ. Math. I.H.E.S.*, 31:59–64, 1966.

[Kon95] Maxim Kontsevich. Lecture at orsay. 1995.

[Loo00] Eduard Looijenga. Motivic measures. preprint, `math.AG/0006220`, 2000.

[Mus01] Mircea Mustaţă. Singularities of Pairs via Jet Schemes. preprint, `math.AG/0102201`, 2001.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO, TOKYO, 153-8914, JAPAN

E-mail address: `t-yasuda@ms.u-tokyo.ac.jp`